

Assessment of Autoimmune Responses Associated with Asbestos Exposure in Libby MT

Jean C. Pfau, Jami J. Sentissi, Greg Weller,
and Elizabeth A. Putnam

doi:10.1289/ehp.7431 (available at <http://dx.doi.org/>)

Online 30 September 2004



Assessment of Autoimmune Responses Associated with Asbestos Exposure in Libby MT

Jean C. Pfau, Jami J. Sentissi, Greg Weller, and Elizabeth A. Putnam
Center for Environmental Health Sciences
Department of Biomedical & Pharmaceutical Sciences
University of Montana, Missoula MT 59812.

Address for correspondence:

Jean C. Pfau, Ph.D.
Center for Environmental Health Sciences
Department of Biomedical and Pharmaceutical Sciences
Skaggs 154
University of Montana
Missoula MT 59812
E-mail: jpfau@selway.umt.edu
Phone: 406-243-4529
FAX: 406-243-2807

Running title: Autoimmune responses with asbestos exposure

Key words: Asbestos, ANA, environmental autoimmunity, immunotoxicology

Acknowledgements

Sincere appreciation is extended to subjects, families, and the CARD Clinic in Libby MT; to Raymond Hamilton (CEHS biostatistics), Curtis Noonan (CEHS epidemiology), and Marvin Fritzler, University of Calgary (Rheumatology) for their expertise; to Theodore Larson, CDC/ATSDR, for TAR data. This work was supported by NIH grants ES-04804 (JP), ES-11676 (EP), and NRSA Fellowship ES-11249 (JP). A conflict of interest was not reported.

Abbreviations

AID Autoimmune disease

ANA Antinuclear antibody.

ARD Asbestos related disease

ENA Extractable Nuclear Antigen.

RF Rheumatoid Factor.

Outline

Abstract

Introduction

Materials and Methods

Results

Discussion

References

Tables

Figure Legends

Figures

ABSTRACT

Systemic autoimmune responses are associated with certain environmental exposures, including crystalline particles such as silica. Positive antinuclear antibody (ANA) tests have been reported in small cohorts exposed to asbestos, but many questions remain regarding the prevalence, pattern, and significance of autoantibodies associated with asbestos exposures. The population in Libby, Montana, provides a unique opportunity for such a study, due to both occupational and environmental exposures that have occurred as a result of the mining of asbestos-contaminated vermiculite near the community. As part of a multi-faceted assessment of the impact of asbestos exposures on this population, this study explored the possibility of exacerbated autoimmune responses. Age and sex-matched sets of 50 serum samples from Libby and Missoula MT (unexposed) were tested for ANA on HEp-2 cells using indirect immunofluorescence (IIF). Data included frequency of positive tests, ANA titers, staining patterns, and scored fluorescence intensity, all against known controls. Serum IgA, rheumatoid factor (RF), and antibodies to extractable nuclear antigen (ENA) were also tested. The Libby samples showed significantly higher frequency of positive ANA and ENA tests, increased mean fluorescence intensity and titers of the ANAs, and higher serum IgA, compared to Missoula samples. In the Libby samples, positive correlations were found between ANA titers and both lung disease severity and extent of exposure. The results support the hypothesis that asbestos exposure is associated with autoimmune responses, and suggests that a relationship exists between those responses and asbestos-related disease processes.